

## Preface

This textbook is intended for use by students of physics, physical chemistry, and theoretical chemistry. The reader is presumed to have a basic knowledge of atomic and quantum physics at the level provided, for example, by the first few chapters in our book *The Physics of Atoms and Quanta*. The student of physics will find here material which should be included in the basic education of every physicist. This book should furthermore allow students to acquire an appreciation of the breadth and variety within the field of molecular physics and its future as a fascinating area of research.

For the student of chemistry, the concepts introduced in this book will provide a theoretical framework for that entire field of study. With the help of these concepts, it is at least in principle possible to reduce the enormous body of empirical chemical knowledge to a few basic principles: those of quantum mechanics. In addition, modern physical methods whose fundamentals are introduced here are becoming increasingly important in chemistry and now represent indispensable tools for the chemist. As examples, we might mention the structural analysis of complex organic compounds, spectroscopic investigation of very rapid reaction processes or, as a practical application, the remote detection of pollutants in the air.

April 1995

Walter Olthoff  
Program Chair  
ECOOP'95

# Organization

ECOOP'95 is organized by the department of Computer Science, University of Århus and AITO (association Internationale pour les Technologies Object) in cooperation with ACM/SIGPLAN.

## Executive Committee

Conference Chair: Ole Lehrmann Madsen (Århus University, DK)  
Program Chair: Walter Olthoff (DFKI GmbH, Germany)  
Organizing Chair: Jørgen Lindskov Knudsen (Århus University, DK)  
Tutorials: Birger Møller-Pedersen  
(Norwegian Computing Center, Norway)  
Workshops: Eric Jul (University of Copenhagen, Denmark)

## Program Committee

Conference Chair: Ole Lehrmann Madsen (Århus University, DK)  
Program Chair: Walter Olthoff (DFKI GmbH, Germany)  
Organizing Chair: Jørgen Lindskov Knudsen (Århus University, DK)  
Tutorials: Birger Møller-Pedersen  
(Norwegian Computing Center, Norway)  
Workshops: Eric Jul (University of Copenhagen, Denmark)

## Referees

V. Andreev	P.C. Bosetti	Del Buono
Bärwolff	Braunschweig	P. Dingus
E. Barrelet	F.W. Büsser	H. Duhm
H.P. Beck	T. Carli	J. Ebert
G. Bernardi	A.B. Clegg	S. Eichenberger
E. Binder	G. Cozzika	and others not included
P.C. Bosetti	S. Dagoret	in this sample...

## Sponsoring Institutions

Bernauer-Budiman Inc., Reading, Mass.  
The Hofmann-International Company, San Louis Obispo, Cal.  
Kramer Industries, Heidelberg, Germany

# Table of Contents

## Hamiltonian Mechanics

Hamiltonian Mechanics unter besonderer Berücksichtigung der höheren Lehranstalten .....	1
<i>Ivar Ekeland (Princeton University), Roger Temam (Université de Paris-Sud), Jeffrey Dean, David Grove, Craig Chambers (Università di Geova), Kim B. Bruce (Stanford University), Elisa Bertino (Digita Research Center)</i>	
Hamiltonian Mechanics2 .....	7
<i>Ivar Ekeland, Roger Temam</i>	
<b>Author Index</b> .....	13
<b>Subject Index</b> .....	17



# Hamiltonian Mechanics unter besonderer Berücksichtigung der höheren Lehranstalten

Ivar Ekeland<sup>1</sup>, Roger Teman<sup>2</sup>, Jeffrey Dean, David Grove, Craig Chambers, Kim B. Bruce, and Elsa Bertino

<sup>1</sup> Princeton University, Princeton NJ 08544, USA,  
I.Ekeland@princeton.edu,

WWW home page: <http://users/~iekeland/web/welcome.html>

<sup>2</sup> Université de Paris-Sud, Laboratoire d'Analyse Numérique, Bâtiment 425,  
F-91405 Orsay Cedex, France

**Abstract.** The abstract should summarize the contents of the paper using at least 70 and at most 150 words. It will be set in 9-point font size and be inset 1.0 cm from the right and left margins. There will be two blank lines before and after the Abstract. . . .

## 1 Fixed-Period Problems: The Sublinear Case

With this chapter, the preliminaries are over, and we begin the search for periodic solutions to Hamiltonian systems. All this will be done in the convex case; that is, we shall study the boundary-value problem

$$\begin{aligned}\dot{x} &= JH'(t, x) \\ x(0) &= x(T)\end{aligned}$$

with  $H(t, \cdot)$  a convex function of  $x$ , going to  $+\infty$  when  $\|x\| \rightarrow \infty$ .

### 1.1 Autonomous Systems

In this section, we will consider the case when the Hamiltonian  $H(x)$  is autonomous. For the sake of simplicity, we shall also assume that it is  $C^1$ .

We shall first consider the question of nontriviality, within the general framework of  $(A_\infty, B_\infty)$ -subquadratic Hamiltonians. In the second subsection, we shall look into the special case when  $H$  is  $(0, b_\infty)$ -subquadratic, and we shall try to derive additional information.

**The General Case: Nontriviality.** We assume that  $H$  is  $(A_\infty, B_\infty)$ -subquadratic at infinity, for some constant symmetric matrices  $A_\infty$  and  $B_\infty$ , with  $B_\infty - A_\infty$  positive definite. Set:

$$\gamma := \text{smallest eigenvalue of } B_\infty - A_\infty \tag{1}$$

$$\lambda := \text{largest negative eigenvalue of } J \frac{d}{dt} + A_\infty . \tag{2}$$

**Fig. 1.** This is the caption of the figure displaying a white eagle and a white horse on a snow field

Theorem 1 tells us that if  $\lambda + \gamma < 0$ , the boundary-value problem:

$$\begin{aligned} \dot{x} &= JH'(x) \\ x(0) &= x(T) \end{aligned} \quad (3)$$

has at least one solution  $\bar{x}$ , which is found by minimizing the dual action functional:

$$\psi(u) = \int_0^T \left[ \frac{1}{2} (A_o^{-1}u, u) + N^*(-u) \right] dt \quad (4)$$

on the range of  $A$ , which is a subspace  $R(A)_L^2$  with finite codimension. Here

$$N(x) := H(x) - \frac{1}{2} (A_\infty x, x) \quad (5)$$

is a convex function, and

$$N(x) \leq \frac{1}{2} ((B_\infty - A_\infty)x, x) + c \quad \forall x . \quad (6)$$

**Proposition 1.** Assume  $H'(0) = 0$  and  $H(0) = 0$ . Set:

$$\delta := \liminf_{x \rightarrow 0} 2N(x) \|x\|^{-2} . \quad (7)$$

If  $\gamma < -\lambda < \delta$ , the solution  $\bar{u}$  is non-zero:

$$\bar{x}(t) \neq 0 \quad \forall t . \quad (8)$$

*Proof.* Condition (7) means that, for every  $\delta' > \delta$ , there is some  $\varepsilon > 0$  such that

$$\|x\| \leq \varepsilon \Rightarrow N(x) \leq \frac{\delta'}{2} \|x\|^2 . \quad (9)$$

It is an exercise in convex analysis, into which we shall not go, to show that this implies that there is an  $\eta > 0$  such that

$$f \|x\| \leq \eta \Rightarrow N^*(y) \leq \frac{1}{2\delta'} \|y\|^2 . \quad (10)$$

Since  $u_1$  is a smooth function, we will have  $\|hu_1\|_\infty \leq \eta$  for  $h$  small enough, and inequality (10) will hold, yielding thereby:

$$\psi(hu_1) \leq \frac{h^2}{2} \frac{1}{\lambda} \|u_1\|_2^2 + \frac{h^2}{2} \frac{1}{\delta'} \|u_1\|^2 . \quad (11)$$

If we choose  $\delta'$  close enough to  $\delta$ , the quantity  $(\frac{1}{\lambda} + \frac{1}{\delta'})$  will be negative, and we end up with

$$\psi(hu_1) < 0 \quad \text{for } h \neq 0 \text{ small} . \quad (12)$$

On the other hand, we check directly that  $\psi(0) = 0$ . This shows that 0 cannot be a minimizer of  $\psi$ , not even a local one. So  $\bar{u} \neq 0$  and  $\bar{u} \neq \Lambda_o^{-1}(0) = 0$ .  $\square$

**Corollary 1.** *Assume  $H$  is  $C^2$  and  $(a_\infty, b_\infty)$ -subquadratic at infinity. Let  $\xi_1, \dots, \xi_N$  be the equilibria, that is, the solutions of  $H'(\xi) = 0$ . Denote by  $\omega_k$  the smallest eigenvalue of  $H''(\xi_k)$ , and set:*

$$\omega := \text{Min} \{ \omega_1, \dots, \omega_k \} . \quad (13)$$

If:

$$\frac{T}{2\pi} b_\infty < -E \left[ -\frac{T}{2\pi} a_\infty \right] < \frac{T}{2\pi} \omega \quad (14)$$

then minimization of  $\psi$  yields a non-constant  $T$ -periodic solution  $\bar{x}$ .

We recall once more that by the integer part  $E[\alpha]$  of  $\alpha \in \mathbb{R}$ , we mean the  $a \in \mathbb{Z}$  such that  $a < \alpha \leq a + 1$ . For instance, if we take  $a_\infty = 0$ , Corollary 2 tells us that  $\bar{x}$  exists and is non-constant provided that:

$$\frac{T}{2\pi} b_\infty < 1 < \frac{T}{2\pi} \quad (15)$$

or

$$T \in \left( \frac{2\pi}{\omega}, \frac{2\pi}{b_\infty} \right) . \quad (16)$$

*Proof.* The spectrum of  $A$  is  $\frac{2\pi}{T} \mathbb{Z} + a_\infty$ . The largest negative eigenvalue  $\lambda$  is given by  $\frac{2\pi}{T} k_o + a_\infty$ , where

$$\frac{2\pi}{T} k_o + a_\infty < 0 \leq \frac{2\pi}{T} (k_o + 1) + a_\infty . \quad (17)$$

Hence:

$$k_o = E \left[ -\frac{T}{2\pi} a_\infty \right] . \quad (18)$$

The condition  $\gamma < -\lambda < \delta$  now becomes:

$$b_\infty - a_\infty < -\frac{2\pi}{T} k_o - a_\infty < \omega - a_\infty \quad (19)$$

which is precisely condition (14).  $\square$

**Table 1.** This is the example table taken out of *The T<sub>E</sub>Xbook*, p. 246

Year	World population
8000 B.C.	5,000,000
50 A.D.	200,000,000
1650 A.D.	500,000,000
1945 A.D.	2,300,000,000
1980 A.D.	4,400,000,000

**Lemma 1.** *Assume that  $H$  is  $C^2$  on  $\mathbb{R}^{2n} \setminus \{0\}$  and that  $H''(x)$  is non-degenerate for any  $x \neq 0$ . Then any local minimizer  $\tilde{x}$  of  $\psi$  has minimal period  $T$ .*

*Proof.* We know that  $\tilde{x}$ , or  $\tilde{x} + \xi$  for some constant  $\xi \in \mathbb{R}^{2n}$ , is a  $T$ -periodic solution of the Hamiltonian system:

$$\dot{x} = JH'(x) . \quad (20)$$

There is no loss of generality in taking  $\xi = 0$ . So  $\psi(x) \geq \psi(\tilde{x})$  for all  $\tilde{x}$  in some neighbourhood of  $x$  in  $W^{1,2}(\mathbb{R}/T\mathbb{Z}; \mathbb{R}^{2n})$ .

But this index is precisely the index  $i_T(\tilde{x})$  of the  $T$ -periodic solution  $\tilde{x}$  over the interval  $(0, T)$ , as defined in Sect. 2.6. So

$$i_T(\tilde{x}) = 0 . \quad (21)$$

Now if  $\tilde{x}$  has a lower period,  $T/k$  say, we would have, by Corollary 31:

$$i_T(\tilde{x}) = i_{kT/k}(\tilde{x}) \geq ki_{T/k}(\tilde{x}) + k - 1 \geq k - 1 \geq 1 . \quad (22)$$

This would contradict (21), and thus cannot happen.  $\square$

*Notes and Comments.* The results in this section are a refined version of [1]; the minimality result of Proposition 14 was the first of its kind.

To understand the nontriviality conditions, such as the one in formula (16), one may think of a one-parameter family  $x_T$ ,  $T \in (2\pi\omega^{-1}, 2\pi b_\infty^{-1})$  of periodic solutions,  $x_T(0) = x_T(T)$ , with  $x_T$  going away to infinity when  $T \rightarrow 2\pi\omega^{-1}$ , which is the period of the linearized system at 0.

**Theorem 1 (Ghoussoub-Preiss).** *Assume  $H(t, x)$  is  $(0, \varepsilon)$ -subquadratic at infinity for all  $\varepsilon > 0$ , and  $T$ -periodic in  $t$*

$$H(t, \cdot) \quad \text{is convex} \quad \forall t \quad (23)$$

$$H(\cdot, x) \quad \text{is } T\text{-periodic} \quad \forall x \quad (24)$$

$$H(t, x) \geq n(\|x\|) \quad \text{with } n(s)s^{-1} \rightarrow \infty \text{ as } s \rightarrow \infty \quad (25)$$

$$\forall \varepsilon > 0, \quad \exists c : H(t, x) \leq \frac{\varepsilon}{2} \|x\|^2 + c . \quad (26)$$

Assume also that  $H$  is  $C^2$ , and  $H''(t, x)$  is positive definite everywhere. Then there is a sequence  $x_k$ ,  $k \in \mathbb{N}$ , of  $kT$ -periodic solutions of the system

$$\dot{x} = JH'(t, x) \quad (27)$$

such that, for every  $k \in \mathbb{N}$ , there is some  $p_o \in \mathbb{N}$  with:

$$p \geq p_o \Rightarrow x_{pk} \neq x_k . \quad (28)$$

□

*Example 1 (External forcing).* Consider the system:

$$\dot{x} = JH'(x) + f(t) \quad (29)$$

where the Hamiltonian  $H$  is  $(0, b_\infty)$ -subquadratic, and the forcing term is a distribution on the circle:

$$f = \frac{d}{dt}F + f_o \quad \text{with } F \in L^2(\mathbb{R}/T\mathbb{Z}; \mathbb{R}^{2n}) , \quad (30)$$

where  $f_o := T^{-1} \int_0^T f(t)dt$ . For instance,

$$f(t) = \sum_{k \in \mathbb{N}} \delta_k \xi , \quad (31)$$

where  $\delta_k$  is the Dirac mass at  $t = k$  and  $\xi \in \mathbb{R}^{2n}$  is a constant, fits the prescription. This means that the system  $\dot{x} = JH'(x)$  is being excited by a series of identical shocks at interval  $T$ .

**Definition 1.** Let  $A_\infty(t)$  and  $B_\infty(t)$  be symmetric operators in  $\mathbb{R}^{2n}$ , depending continuously on  $t \in [0, T]$ , such that  $A_\infty(t) \leq B_\infty(t)$  for all  $t$ .

A Borelian function  $H : [0, T] \times \mathbb{R}^{2n} \rightarrow \mathbb{R}$  is called  $(A_\infty, B_\infty)$ -subquadratic at infinity if there exists a function  $N(t, x)$  such that:

$$H(t, x) = \frac{1}{2} (A_\infty(t)x, x) + N(t, x) \quad (32)$$

$$\forall t , \quad N(t, x) \quad \text{is convex with respect to } x \quad (33)$$

$$N(t, x) \geq n(\|x\|) \quad \text{with } n(s)s^{-1} \rightarrow +\infty \text{ as } s \rightarrow +\infty \quad (34)$$

$$\exists c \in \mathbb{R} : \quad H(t, x) \leq \frac{1}{2} (B_\infty(t)x, x) + c \quad \forall x . \quad (35)$$

If  $A_\infty(t) = a_\infty I$  and  $B_\infty(t) = b_\infty I$ , with  $a_\infty \leq b_\infty \in \mathbb{R}$ , we shall say that  $H$  is  $(a_\infty, b_\infty)$ -subquadratic at infinity. As an example, the function  $\|x\|^\alpha$ , with  $1 \leq \alpha < 2$ , is  $(0, \varepsilon)$ -subquadratic at infinity for every  $\varepsilon > 0$ . Similarly, the Hamiltonian

$$H(t, x) = \frac{1}{2}k \|k\|^2 + \|x\|^\alpha \quad (36)$$

is  $(k, k + \varepsilon)$ -subquadratic for every  $\varepsilon > 0$ . Note that, if  $k < 0$ , it is not convex.

*Notes and Comments.* The first results on subharmonics were obtained by Rabinowitz in [5], who showed the existence of infinitely many subharmonics both in the subquadratic and superquadratic case, with suitable growth conditions on  $H'$ . Again the duality approach enabled Clarke and Ekeland in [2] to treat the same problem in the convex-subquadratic case, with growth conditions on  $H$  only.

Recently, Michalek and Tarantello (see [3] and [4]) have obtained lower bound on the number of subharmonics of period  $kT$ , based on symmetry considerations and on pinching estimates, as in Sect. 5.2 of this article.

## References

1. Clarke, F., Ekeland, I.: Nonlinear oscillations and boundary-value problems for Hamiltonian systems. *Arch. Rat. Mech. Anal.* **78** (1982) 315–333
2. Clarke, F., Ekeland, I.: Solutions périodiques, du période donnée, des équations hamiltoniennes. *Note CRAS Paris* **287** (1978) 1013–1015
3. Michalek, R., Tarantello, G.: Subharmonic solutions with prescribed minimal period for nonautonomous Hamiltonian systems. *J. Diff. Eq.* **72** (1988) 28–55
4. Tarantello, G.: Subharmonic solutions for Hamiltonian systems via a  $\mathbb{Z}_p$  pseudoinde index theory. *Annali di Matematica Pura* (to appear)
5. Rabinowitz, P.: On subharmonic solutions of a Hamiltonian system. *Comm. Pure Appl. Math.* **33** (1980) 609–633

# Hamiltonian Mechanics2

Ivar Ekeland<sup>1</sup> and Roger Temam<sup>2</sup>

<sup>1</sup> Princeton University, Princeton NJ 08544, USA

<sup>2</sup> Université de Paris-Sud, Laboratoire d'Analyse Numérique, Bâtiment 425,  
F-91405 Orsay Cedex, France

**Abstract.** The abstract should summarize the contents of the paper using at least 70 and at most 150 words. It will be set in 9-point font size and be inset 1.0 cm from the right and left margins. There will be two blank lines before and after the Abstract. . .

## 1 Fixed-Period Problems: The Sublinear Case

With this chapter, the preliminaries are over, and we begin the search for periodic solutions to Hamiltonian systems. All this will be done in the convex case; that is, we shall study the boundary-value problem

$$\begin{aligned}\dot{x} &= JH'(t, x) \\ x(0) &= x(T)\end{aligned}$$

with  $H(t, \cdot)$  a convex function of  $x$ , going to  $+\infty$  when  $\|x\| \rightarrow \infty$ .

### 1.1 Autonomous Systems

In this section, we will consider the case when the Hamiltonian  $H(x)$  is autonomous. For the sake of simplicity, we shall also assume that it is  $C^1$ .

We shall first consider the question of nontriviality, within the general framework of  $(A_\infty, B_\infty)$ -subquadratic Hamiltonians. In the second subsection, we shall look into the special case when  $H$  is  $(0, b_\infty)$ -subquadratic, and we shall try to derive additional information.

**The General Case: Nontriviality.** We assume that  $H$  is  $(A_\infty, B_\infty)$ -subquadratic at infinity, for some constant symmetric matrices  $A_\infty$  and  $B_\infty$ , with  $B_\infty - A_\infty$  positive definite. Set:

$$\gamma := \text{smallest eigenvalue of } B_\infty - A_\infty \tag{1}$$

$$\lambda := \text{largest negative eigenvalue of } J \frac{d}{dt} + A_\infty . \tag{2}$$

Theorem 21 tells us that if  $\lambda + \gamma < 0$ , the boundary-value problem:

$$\begin{aligned}\dot{x} &= JH'(x) \\ x(0) &= x(T)\end{aligned} \tag{3}$$

**Fig. 1.** This is the caption of the figure displaying a white eagle and a white horse on a snow field

has at least one solution  $\bar{x}$ , which is found by minimizing the dual action functional:

$$\psi(u) = \int_0^T \left[ \frac{1}{2} (A_o^{-1}u, u) + N^*(-u) \right] dt \quad (4)$$

on the range of  $A$ , which is a subspace  $R(A)_L^2$  with finite codimension. Here

$$N(x) := H(x) - \frac{1}{2} (A_\infty x, x) \quad (5)$$

is a convex function, and

$$N(x) \leq \frac{1}{2} ((B_\infty - A_\infty)x, x) + c \quad \forall x . \quad (6)$$

**Proposition 1.** Assume  $H'(0) = 0$  and  $H(0) = 0$ . Set:

$$\delta := \liminf_{x \rightarrow 0} 2N(x) \|x\|^{-2} . \quad (7)$$

If  $\gamma < -\lambda < \delta$ , the solution  $\bar{u}$  is non-zero:

$$\bar{x}(t) \neq 0 \quad \forall t . \quad (8)$$

*Proof.* Condition (7) means that, for every  $\delta' > \delta$ , there is some  $\varepsilon > 0$  such that

$$\|x\| \leq \varepsilon \Rightarrow N(x) \leq \frac{\delta'}{2} \|x\|^2 . \quad (9)$$

It is an exercise in convex analysis, into which we shall not go, to show that this implies that there is an  $\eta > 0$  such that

$$f \|x\| \leq \eta \Rightarrow N^*(y) \leq \frac{1}{2\delta'} \|y\|^2 . \quad (10)$$

Since  $u_1$  is a smooth function, we will have  $\|hu_1\|_\infty \leq \eta$  for  $h$  small enough, and inequality (10) will hold, yielding thereby:

$$\psi(hu_1) \leq \frac{h^2}{2} \frac{1}{\lambda} \|u_1\|_2^2 + \frac{h^2}{2} \frac{1}{\delta'} \|u_1\|^2 . \quad (11)$$

If we choose  $\delta'$  close enough to  $\delta$ , the quantity  $(\frac{1}{\lambda} + \frac{1}{\delta'})$  will be negative, and we end up with

$$\psi(hu_1) < 0 \quad \text{for } h \neq 0 \text{ small} . \quad (12)$$

On the other hand, we check directly that  $\psi(0) = 0$ . This shows that 0 cannot be a minimizer of  $\psi$ , not even a local one. So  $\bar{u} \neq 0$  and  $\bar{u} \neq \Lambda_o^{-1}(0) = 0$ .  $\square$

**Corollary 1.** *Assume  $H$  is  $C^2$  and  $(a_\infty, b_\infty)$ -subquadratic at infinity. Let  $\xi_1, \dots, \xi_N$  be the equilibria, that is, the solutions of  $H'(\xi) = 0$ . Denote by  $\omega_k$  the smallest eigenvalue of  $H''(\xi_k)$ , and set:*

$$\omega := \text{Min} \{ \omega_1, \dots, \omega_k \} . \quad (13)$$

If:

$$\frac{T}{2\pi} b_\infty < -E \left[ -\frac{T}{2\pi} a_\infty \right] < \frac{T}{2\pi} \omega \quad (14)$$

then minimization of  $\psi$  yields a non-constant  $T$ -periodic solution  $\bar{x}$ .

We recall once more that by the integer part  $E[\alpha]$  of  $\alpha \in \mathbb{R}$ , we mean the  $a \in \mathbb{Z}$  such that  $a < \alpha \leq a + 1$ . For instance, if we take  $a_\infty = 0$ , Corollary 2 tells us that  $\bar{x}$  exists and is non-constant provided that:

$$\frac{T}{2\pi} b_\infty < 1 < \frac{T}{2\pi} \quad (15)$$

or

$$T \in \left( \frac{2\pi}{\omega}, \frac{2\pi}{b_\infty} \right) . \quad (16)$$

*Proof.* The spectrum of  $\Lambda$  is  $\frac{2\pi}{T}\mathbb{Z} + a_\infty$ . The largest negative eigenvalue  $\lambda$  is given by  $\frac{2\pi}{T}k_o + a_\infty$ , where

$$\frac{2\pi}{T}k_o + a_\infty < 0 \leq \frac{2\pi}{T}(k_o + 1) + a_\infty . \quad (17)$$

Hence:

$$k_o = E \left[ -\frac{T}{2\pi} a_\infty \right] . \quad (18)$$

The condition  $\gamma < -\lambda < \delta$  now becomes:

$$b_\infty - a_\infty < -\frac{2\pi}{T}k_o - a_\infty < \omega - a_\infty \quad (19)$$

which is precisely condition (14).  $\square$

**Lemma 1.** *Assume that  $H$  is  $C^2$  on  $\mathbb{R}^{2n} \setminus \{0\}$  and that  $H''(x)$  is non-degenerate for any  $x \neq 0$ . Then any local minimizer  $\bar{x}$  of  $\psi$  has minimal period  $T$ .*

**Table 1.** This is the example table taken out of *The T<sub>E</sub>Xbook*, p. 246

Year	World population
8000 B.C.	5,000,000
50 A.D.	200,000,000
1650 A.D.	500,000,000
1945 A.D.	2,300,000,000
1980 A.D.	4,400,000,000

*Proof.* We know that  $\tilde{x}$ , or  $\tilde{x} + \xi$  for some constant  $\xi \in \mathbb{R}^{2n}$ , is a  $T$ -periodic solution of the Hamiltonian system:

$$\dot{x} = JH'(x) . \quad (20)$$

There is no loss of generality in taking  $\xi = 0$ . So  $\psi(x) \geq \psi(\tilde{x})$  for all  $\tilde{x}$  in some neighbourhood of  $x$  in  $W^{1,2}(\mathbb{R}/T\mathbb{Z}; \mathbb{R}^{2n})$ .

But this index is precisely the index  $i_T(\tilde{x})$  of the  $T$ -periodic solution  $\tilde{x}$  over the interval  $(0, T)$ , as defined in Sect. 2.6. So

$$i_T(\tilde{x}) = 0 . \quad (21)$$

Now if  $\tilde{x}$  has a lower period,  $T/k$  say, we would have, by Corollary 31:

$$i_T(\tilde{x}) = i_{kT/k}(\tilde{x}) \geq ki_{T/k}(\tilde{x}) + k - 1 \geq k - 1 \geq 1 . \quad (22)$$

This would contradict (21), and thus cannot happen.  $\square$

*Notes and Comments.* The results in this section are a refined version of 1980; the minimality result of Proposition 14 was the first of its kind.

To understand the nontriviality conditions, such as the one in formula (16), one may think of a one-parameter family  $x_T$ ,  $T \in (2\pi\omega^{-1}, 2\pi b_\infty^{-1})$  of periodic solutions,  $x_T(0) = x_T(T)$ , with  $x_T$  going away to infinity when  $T \rightarrow 2\pi\omega^{-1}$ , which is the period of the linearized system at 0.

**Theorem 1 (Ghoussoub-Preiss).** *Assume  $H(t, x)$  is  $(0, \varepsilon)$ -subquadratic at infinity for all  $\varepsilon > 0$ , and  $T$ -periodic in  $t$*

$$H(t, \cdot) \quad \text{is convex} \quad \forall t \quad (23)$$

$$H(\cdot, x) \quad \text{is } T\text{-periodic} \quad \forall x \quad (24)$$

$$H(t, x) \geq n(\|x\|) \quad \text{with } n(s)s^{-1} \rightarrow \infty \text{ as } s \rightarrow \infty \quad (25)$$

$$\forall \varepsilon > 0, \quad \exists c : H(t, x) \leq \frac{\varepsilon}{2} \|x\|^2 + c . \quad (26)$$

*Assume also that  $H$  is  $C^2$ , and  $H''(t, x)$  is positive definite everywhere. Then there is a sequence  $x_k$ ,  $k \in \mathbb{N}$ , of  $kT$ -periodic solutions of the system*

$$\dot{x} = JH'(t, x) \quad (27)$$

such that, for every  $k \in \mathbb{N}$ , there is some  $p_o \in \mathbb{N}$  with:

$$p \geq p_o \Rightarrow x_{pk} \neq x_k . \quad (28)$$

□

*Example 1 (External forcing).* Consider the system:

$$\dot{x} = JH'(x) + f(t) \quad (29)$$

where the Hamiltonian  $H$  is  $(0, b_\infty)$ -subquadratic, and the forcing term is a distribution on the circle:

$$f = \frac{d}{dt}F + f_o \quad \text{with } F \in L^2(\mathbb{R}/T\mathbb{Z}; \mathbb{R}^{2n}) , \quad (30)$$

where  $f_o := T^{-1} \int_o^T f(t)dt$ . For instance,

$$f(t) = \sum_{k \in \mathbb{N}} \delta_k \xi , \quad (31)$$

where  $\delta_k$  is the Dirac mass at  $t = k$  and  $\xi \in \mathbb{R}^{2n}$  is a constant, fits the prescription. This means that the system  $\dot{x} = JH'(x)$  is being excited by a series of identical shocks at interval  $T$ .

**Definition 1.** Let  $A_\infty(t)$  and  $B_\infty(t)$  be symmetric operators in  $\mathbb{R}^{2n}$ , depending continuously on  $t \in [0, T]$ , such that  $A_\infty(t) \leq B_\infty(t)$  for all  $t$ .

A Borelian function  $H : [0, T] \times \mathbb{R}^{2n} \rightarrow \mathbb{R}$  is called  $(A_\infty, B_\infty)$ -subquadratic at infinity if there exists a function  $N(t, x)$  such that:

$$H(t, x) = \frac{1}{2} (A_\infty(t)x, x) + N(t, x) \quad (32)$$

$$\forall t , \quad N(t, x) \quad \text{is convex with respect to } x \quad (33)$$

$$N(t, x) \geq n(\|x\|) \quad \text{with } n(s)s^{-1} \rightarrow +\infty \text{ as } s \rightarrow +\infty \quad (34)$$

$$\exists c \in \mathbb{R} : \quad H(t, x) \leq \frac{1}{2} (B_\infty(t)x, x) + c \quad \forall x . \quad (35)$$

If  $A_\infty(t) = a_\infty I$  and  $B_\infty(t) = b_\infty I$ , with  $a_\infty \leq b_\infty \in \mathbb{R}$ , we shall say that  $H$  is  $(a_\infty, b_\infty)$ -subquadratic at infinity. As an example, the function  $\|x\|^\alpha$ , with  $1 \leq \alpha < 2$ , is  $(0, \varepsilon)$ -subquadratic at infinity for every  $\varepsilon > 0$ . Similarly, the Hamiltonian

$$H(t, x) = \frac{1}{2}k \|k\|^2 + \|x\|^\alpha \quad (36)$$

is  $(k, k + \varepsilon)$ -subquadratic for every  $\varepsilon > 0$ . Note that, if  $k < 0$ , it is not convex.

*Notes and Comments.* The first results on subharmonics were obtained by Rabinowitz in 1985, who showed the existence of infinitely many subharmonics both in the subquadratic and superquadratic case, with suitable growth conditions on  $H'$ . Again the duality approach enabled Clarke and Ekeland in 1981 to treat the same problem in the convex-subquadratic case, with growth conditions on  $H$  only.

Recently, Michalek and Tarantello (see Michalek, R., Tarantello, G. 1982 and Tarantello, G. 1983) have obtained lower bound on the number of subharmonics of period  $kT$ , based on symmetry considerations and on pinching estimates, as in Sect. 5.2 of this article.

## References

- Clarke, F., Ekeland, I.: Nonlinear oscillations and boundary-value problems for Hamiltonian systems. *Arch. Rat. Mech. Anal.* **78** (1982) 315–333
- Clarke, F., Ekeland, I.: Solutions périodiques, du période donnée, des équations hamiltoniennes. *Note CRAS Paris* **287** (1978) 1013–1015
- Michalek, R., Tarantello, G.: Subharmonic solutions with prescribed minimal period for nonautonomous Hamiltonian systems. *J. Diff. Eq.* **72** (1988) 28–55
- Tarantello, G.: Subharmonic solutions for Hamiltonian systems via a  $\mathbb{Z}_p$  pseudoindex theory. *Annali di Matematica Pura* (to appear)
- Rabinowitz, P.: On subharmonic solutions of a Hamiltonian system. *Comm. Pure Appl. Math.* **33** (1980) 609–633

## Author Index

- Abt I. 7  
Ahmed T. 3  
Andreev V. 24  
Andrieu B. 27  
Arpagaus M. 34
- Babaev A. 25  
Bärwolf A. 33  
Bán J. 17  
Baranov P. 24  
Barrelet E. 28  
Bartel W. 11  
Bassler U. 28  
Beck H.P. 35  
Behrend H.-J. 11  
Berger Ch. 1  
Bergstein H. 1  
Bernardi G. 28  
Bernet R. 34  
Besançon M. 9  
Biddulph P. 22  
Binder E. 11  
Bischoff A. 33  
Blobel V. 13  
Borras K. 8  
Bosetti P.C. 2  
Boudry V. 27  
Brasse F. 11  
Braun U. 2  
Braunschweig A. 1  
Brisson V. 26  
Büngener L. 13  
Bürger J. 11  
Büsser F.W. 13  
Buniatian A. 11,37  
Buschhorn G. 25
- Campbell A.J. 1  
Carli T. 25  
Charles F. 28  
Clarke D. 5  
Clegg A.B. 18  
Colombo M. 8  
Courau A. 26  
Coutures Ch. 9
- Cozzika G. 9  
Criegee L. 11  
Cvach J. 27
- Dagoret S. 28  
Dainton J.B. 19  
Dann A.W.E. 22  
Dau W.D. 16  
Deffur E. 11  
Delcourt B. 26  
Buono Del A. 28  
Devel M. 26  
De Roeck A. 11  
Dingus P. 27  
Dollfus C. 35  
Dreis H.B. 2  
Drescher A. 8  
Düllmann D. 13  
Dünger O. 13  
Duhm H. 12
- Ebbinghaus R. 8  
Eberle M. 12  
Ebert J. 32  
Ebert T.R. 19  
Efremenko V. 23  
Egli S. 35  
Eichenberger S. 35  
Eichler R. 34  
Eisenhandler E. 20  
Ellis N.N. 3  
Ellison R.J. 22  
Elsen E. 11  
Evrard E. 4
- Favart L. 4  
Feeken D. 13  
Felst R. 11  
Feltesse A. 9  
Fensome I.F. 3  
Ferrarotto F. 31  
Flamm K. 11  
Flauger W. 11  
Flieser M. 25  
Flügge G. 2

- Fomenko A. 24  
 Fominykh B. 23  
 Formánek J. 30  
 Foster J.M. 22  
 Franke G. 11  
 Fretwurst E. 12  
  
 Gabathuler E. 19  
 Gamedinger K. 25  
 Garvey J. 3  
 Gayler J. 11  
 Gellrich A. 13  
 Gennis M. 11  
 Genzel H. 1  
 Godfrey L. 7  
 Goerlach U. 11  
 Goerlich L. 6  
 Gogitidze N. 24  
 Goodall A.M. 19  
 Gorelov I. 23  
 Goritchev P. 23  
 Grab C. 34  
 Grässler R. 2  
 Greenshaw T. 19  
 Greif H. 25  
 Grindhammer G. 25  
  
 Haack J. 33  
 Haidt D. 11  
 Hamon O. 28  
 Handschuh D. 11  
 Hanlon E.M. 18  
 Hapke M. 11  
 Harjes J. 11  
 Haydar R. 26  
 Haynes W.J. 5  
 Hedberg V. 21  
 Heinzelmann G. 13  
 Henderson R.C.W. 18  
 Henschel H. 33  
 Herynek I. 29  
 Hildesheim W. 11  
 Hill P. 11  
 Hilton C.D. 22  
 Hoeger K.C. 22  
 Huet Ph. 4  
 Hufnagel H. 14  
 Huot N. 28  
  
 Itterbeck H. 1  
  
 Jabiol M.-A. 9  
 Jacholkowska A. 26  
 Jacobsson C. 21  
 Jansen T. 11  
 Jönsson L. 21  
 Johannsen A. 13  
 Johnson D.P. 4  
 Jung H. 2  
  
 Kalmus P.I.P. 20  
 Kasarian S. 11  
 Kaschowitz R. 2  
 Kathage U. 16  
 Kaufmann H. 33  
 Kenyon I.R. 3  
 Kermiche S. 26  
 Kiesling C. 25  
 Klein M. 33  
 Kleinwort C. 13  
 Knies G. 11  
 Ko W. 7  
 Köhler T. 1  
 Kolanoski H. 8  
 Kole F. 7  
 Kolya S.D. 22  
 Korbel V. 11  
 Korn M. 8  
 Kostka P. 33  
 Kotelnikov S.K. 24  
 Krehbiel H. 11  
 Krücker D. 2  
 Krüger U. 11  
 Kubenka J.P. 25  
 Kuhlen M. 25  
 Kurča T. 17  
 Kurzhöfer J. 8  
 Kuznik B. 32  
  
 Lamarche F. 27  
 Lander R. 7  
 Landon M.P.J. 20  
 Lange W. 33  
 Lanius P. 25  
 Laporte J.F. 9  
 Lebedev A. 24  
 Leuschner A. 11  
 Levonian S. 11,24  
 Lewin D. 11  
 Ley Ch. 2  
 Lindner A. 8

Lindström G. 12  
 Linsel F. 11  
 Lipinski J. 13  
 Loch P. 11  
 Lohmander H. 21  
 Lopez G.C. 20  
  
 Magnussen N. 32  
 Mani S. 7  
 Marage P. 4  
 Marshall R. 22  
 Martens J. 32  
 Martin A.@ 19  
 Martyn H.-U. 1  
 Martyniak J. 6  
 Masson S. 2  
 Mavroidis A. 20  
 McMahon S.J. 19  
 Mehta A. 22  
 Meier K. 15  
 Mercer D. 22  
 Merz T. 11  
 Meyer C.A. 35  
 Meyer H. 32  
 Meyer J. 11  
 Mikocki S. 6,26  
 Milone V. 31  
 Moreau F. 27  
 Moreels J. 4  
 Morris J.V. 5  
 Müller K. 35  
 Murray S.A. 22  
  
 Nagovizin V. 23  
 Naroska B. 13  
 Naumann Th. 33  
 Newton D. 18  
 Neyret D. 28  
 Nguyen A. 28  
 Niebergall F. 13  
 Nisius R. 1  
 Nowak G. 6  
 Nyberg M. 21  
  
 Oberlack H. 25  
 Obrock U. 8  
 Olsson J.E. 11  
 Ould-Saada F. 13  
  
 Pascaud C. 26  
  
 Patel G.D. 19  
 Peppel E. 11  
 Phillips H.T. 3  
 Phillips J.P. 22  
 Pichler Ch. 12  
 Pilgram W. 2  
 Pitzl D. 34  
 Prell S. 11  
 Prosi R. 11  
  
 Rädcl G. 11  
 Raupach F. 1  
 Rauschnabel K. 8  
 Reinshagen S. 11  
 Ribarics P. 25  
 Riech V. 12  
 Riedlberger J. 34  
 Rietz M. 2  
 Robertson S.M. 3  
 Robmann P. 35  
 Roosen R. 4  
 Royon C. 9  
 Rudowicz M. 25  
 Rusakov S. 24  
 Rybicki K. 6  
  
 Sahlmann N. 2  
 Sanchez E. 25  
 Savitsky M. 11  
 Schacht P. 25  
 Schleper P. 14  
 von Schlippe W. 20  
 Schmidt D. 32  
 Schmitz W. 2  
 Schöning A. 11  
 Schröder V. 11  
 Schulz M. 11  
 Schwab B. 14  
 Schwind A. 33  
 Seehausen U. 13  
 Sell R. 11  
 Semenov A. 23  
 Shekelyan V. 23  
 Shooshtari H. 25  
 Shtarkov L.N. 24  
 Siegmon G. 16  
 Siewert U. 16  
 Skillicorn I.O. 10  
 Smirnov P. 24  
 Smith J.R. 7

- Smolik L. 11  
Spitzer H. 13  
Staroba P. 29  
Steenbock M. 13  
Steffen P. 11  
Stella B. 31  
Stephens K. 22  
Stösslein U. 33  
Strachota J. 11  
Straumann U. 35  
Struczinski W. 2
- Taylor R.E. 36,26  
Tchernyshov V. 23  
Thiebaux C. 27  
Thompson G. 20  
Truöl P. 35  
Turnau J. 6
- Urban L. 25  
Usik A. 24
- Valkarova A. 30  
Vallée C. 28  
Van Esch P. 4  
Vartapetian A. 11
- Vazdik Y. 24  
Verrecchia P. 9  
Vick R. 13  
Vogel E. 1
- Wacker K. 8  
Walther A. 8  
Weber G. 13  
Wegner A. 11  
Wellisch H.P. 25  
West L.R. 3  
Willard S. 7  
Winde M. 33  
Winter G.-G. 11  
Wolff Th. 34  
Wright A.E. 22  
Wulff N. 11
- Yiou T.P. 28
- Žáček J. 30  
Zeitnitz C. 12  
Ziaepour H. 26  
Zimmer M. 11  
Zimmermann W. 11

# Subject Index

- Absorption 327
- Absorption of radiation 289–292, 299, 300
- Actinides 244
- Aharonov-Bohm effect 142–146
- Angular momentum 101–112
  - algebraic treatment 391–396
- Angular momentum addition 185–193
- Angular momentum commutation relations 101
- Angular momentum quantization 9–10, 104–106
- Angular momentum states 107, 321, 391–396
- Antiquark 83
- $\alpha$ -rays 101–103
- Atomic theory 8–10, 219–249, 327
- Average value
  - (*see also* Expectation value) 15–16, 25, 34, 37, 357
  
- Baker-Hausdorff formula 23
- Balmer formula 8
- Balmer series 125
- Baryon 220, 224
- Basis 98
- Basis system 164, 376
- Bell inequality 379–381, 382
- Bessel functions 201, 313, 337
  - spherical 304–306, 309, 313–314, 322
- Bound state 73–74, 78–79, 116–118, 202, 267, 273, 306, 348, 351
- Boundary conditions 59, 70
- Bra 159
- Breit-Wigner formula 80, 84, 332
  
- Brillouin-Wigner perturbation theory 203
  
- Cathode rays 8
- Causality 357–359
- Center-of-mass frame 232, 274, 338
- Central potential 113–135, 303–314
- Centrifugal potential 115–116, 323
- Characteristic function 33
- Clebsch-Gordan coefficients 191–193
- Cold emission 88
- Combination principle, Ritz’s 124
- Commutation relations 27, 44, 353, 391
- Commutator 21–22, 27, 44, 344
- Compatibility of measurements 99
- Complete orthonormal set 31, 40, 160, 360
- Complete orthonormal system, *see* Complete orthonormal set
- Complete orthonormal set
- Complete set of observables, *see* Complete set of operators
  
- Eigenfunction 34, 46, 344–346
  - radial 321
  - calculation 322–324
- EPR argument 377–378
- Exchange term 228, 231, 237, 241, 268, 272
  
- $f$ -sum rule 302
- Fermi energy 223
  
- H<sub>2</sub><sup>+</sup> molecule 26
- Half-life 65
- Holzwarth energies 68