

# The Title of a TU Wien Report

Dr. Samuel Author

The Date

---

## I. The First Part

---

---

# 1 Sample Mathematics and Text

---

## 1.1 In-line and Displayed Mathematics

The expression  $\sum_{i=1}^{\infty} a_i$  is in-line mathematics, while the numbered equation

$$\sum_{i=1}^{\infty} a_i \tag{1.1}$$

is displayed and automatically numbered as equation 1.1.

Let  $H$  be a Hilbert space,  $C$  be a closed bounded convex subset of  $H$ ,  $T$  a nonexpansive self map of  $C$ . Suppose that as  $n \rightarrow \infty$ ,  $a_{n,k} \rightarrow 0$  for each  $k$ , and  $\gamma_n = \sum_{k=0}^{\infty} (a_{n,k+1} - a_{n,k})^+ \rightarrow 0$ . Then for each  $x$  in  $C$ ,  $A_n x = \sum_{k=0}^{\infty} a_{n,k} T^k x$  converges weakly to a fixed point of  $T$  [1].

Two sets of L<sup>A</sup>T<sub>E</sub>X parameters govern mathematical displays.<sup>1</sup> The spacing above and below a display depends on whether the lines above or below are short or long, as shown in the following examples.

A short line above:

$$x^2 + y^2 = z^2$$

and a short line below.

A long line above may depend on your margins

$$\sin^2 \theta + \cos^2 \theta = 1$$

as will a long line below. This line is long enough to illustrate the spacing for mathematical displays, regardless of the margins.

## 1.2 Mathematics in Section Heads $\int_{\alpha}^{\beta} \ln t dt$

Mathematics can appear in section heads. Note that mathematics in section heads may cause difficulties in typesetting styles with running headers or table of contents entries.

## 1.3 Theorems, Lemmata, and Other Theorem-like Environments

A number of theorem-like environments is available. The following lemma is a well-known fact on differentiation of asymptotic expansions of analytic functions.

**Lemma 1** *Let  $f(z)$  be an analytic function in  $\mathbb{C}_+$ . If  $f(z)$  admits the representation*

$$f(z) = a_0 + \frac{a_1}{z} + o\left(\frac{1}{z}\right),$$

---

<sup>1</sup> L<sup>A</sup>T<sub>E</sub>X automatically selects the spacing depending on the surrounding line lengths.

for  $z \rightarrow \infty$  inside a cone  $\Gamma_\varepsilon = \{z \in \mathbb{C}_+ : 0 < \varepsilon \leq \arg z \leq \pi - \varepsilon\}$  then

$$a_1 = -\lim_{z \rightarrow \infty} z^2 f'(z), \quad z \in \Gamma_\varepsilon. \quad (1.2)$$

**Proof.** Change  $z$  for  $1/z$ . Then  $\Gamma_\varepsilon \rightarrow \bar{\Gamma}_\varepsilon = \{z \in \mathbb{C}_- : \bar{z} \in \Gamma_\varepsilon\}$  and

$$f(1/z) = a_0 + a_1 z + o(z). \quad (1.3)$$

Fix  $z \in \bar{\Gamma}_\varepsilon$ , and let  $C_r(z) = \{\lambda \in \mathbb{C}_- : |\lambda - z| = r\}$  be a circle with radius  $r = |z| \sin \varepsilon/2$ . It follows from (1.3) that

$$\frac{1}{2\pi i} \int_{C_r(z)} \frac{f(\lambda) d\lambda}{(\lambda - z)^2} = \sum_{m=0}^1 a_m \frac{1}{2\pi i} \int_{C_r(z)} \frac{(\lambda - z_0)^m d\lambda}{(\lambda - z)^2} + R(z), \quad (1.4)$$

where for the remainder  $R(z)$  we have

$$\begin{aligned} |R(z)| &\leq r^{-1} \max_{\lambda \in C_r(z)} o(|z|) = r^{-1} \max_{\lambda \in C_r(z)} |\lambda| \cdot O(|z| + r) \\ &= \frac{|z| + r}{r} \cdot O(|z| + r) = \frac{1 + \sin \varepsilon}{\sin \varepsilon} \cdot O(|z|). \end{aligned}$$

Therefore  $R(z) \rightarrow 0$  as  $z \rightarrow \infty$ ,  $z \in \bar{\Gamma}_{\varepsilon/2}$ , and hence by the Cauchy theorem (1.4) implies

$$\frac{d}{dz} f(1/z) = a_1 + R(z) \rightarrow a_1, \quad \text{as } z \rightarrow \infty, \quad z \in \bar{\Gamma}_{\varepsilon/2},$$

that implies (1.2) by substituting  $1/z$  back for  $z$ . ■

---

## 2 Features of this Shell

---

### 2.1 Section Headings

Use the Section tag for major sections, and the Subsection tag for subsections.

#### 2.1.1 Subsection

This is some harmless text under a subsection.

#### Subsubsection

This is some harmless text under a subsubsection.

**Subsubsubsection** This is some harmless text under a subsubsubsection.

**Subsubsubsubsection** This is some harmless text under a subsubsubsubsection.

### 2.2 Tags

You can apply the logical markup tag *Emphasized*.

You can apply the visual markup tags **Bold**, *Italics*, Roman, Sans Serif, *Slanted*, SMALL CAPS, and **Typewriter**.

You can apply the special, mathematics only, tags BLACKBOARD BOLD, *CALLIGRAPHIC*, and  $\text{fraktur}$ . Note that blackboard bold and calligraphic are correct only when applied to uppercase letters A through Z.

You can apply the size tags tiny, scriptsize, footnotesize, small, normalsize, large, Large, LARGE, huge and Huge.

This is a Body Math paragraph. Each time you press the Enter key, Scientific WorkPlace switches to mathematics mode. This is convenient for carrying out “scratchpad” computations.

Following is a group of paragraphs marked as Body Quote. This environment is appropriate for a short quotation or a sequence of short quotations.

The only thing we have to fear is fear itself. *Franklin D. Roosevelt*, Mar. 4, 1933

Ask not what your country can do for you; ask what you can do for your country. *John F. Kennedy*, Jan. 20, 1961

There is nothing wrong with America that cannot be cured by what is right with America. *William J. “Bill” Clinton*, Jan. 21, 1993

## 2.3 List Environments

You can create numbered, bulleted, and description lists using the tag popup at the bottom left of the screen.

1. List item 1
2. List item 2

(a) A list item under a list item.

The typeset style for this level is different than the screen style. The screen shows a lower case alphabetic character followed by a period while the typeset style uses a lower case alphabetic character surrounded by parentheses.

(b) Just another list item under a list item.

i. Third level list item under a list item.

A. Fourth and final level of list items allowed.

- Bullet item 1
- Bullet item 2

– Second level bullet item.

\* Third level bullet item.

· Fourth (and final) level bullet item.

Description List : Each description list item has a term followed by the description of that term. Double click the term box to enter the term, or to change it.

Bunyip : Mythical beast of Australian Aboriginal legends.

---

## Appendix A The First Appendix

---

The appendix fragment is used only once. Subsequent appendices can be created using the Chapter Section/Body Tag.

### A.1 About the Bibliography

Following the text of this article is a short manual bibliography. This sample bibliography has no relationship to the previous text, but it shows sample citations such as [4], [5] and [6]. You can also have multiple citations appear together. Here is an example: [2, 3, 4].

---

## Bibliography

---

- [1] N. Dunford and J. Schwartz, *Functional Analysis*, v. 2, John Wiley and Sons, New York, 1963.
- [2] Harstad, K. and Bellan, J., "Isolated fluid oxygen drop behavior in fluid hydrogen at rocket chamber pressures", *Int. J. Heat Mass Transfer*, 1998a, **41**, 3537-3550
- [3] Harstad, K. and Bellan, J., "The Lewis number under supercritical conditions", *Int. J. Heat Mass Transfer*, in print
- [4] Hirshfelder, J. O., Curtis, C. F. and Bird, R. B., *Molecular Theory of Gases and Liquids*, John Wiley and Sons, Inc., 1964
- [5] Prausnitz, J., Lichtenthaler, R. and de Azevedo, E., *Molecular thermodynamics for fluid-phase equilibrium*, Prentice -Hall, Inc., 1986
- [6] Reid, R. C., Prausnitz, J. M. and Polling, B. E., *The Properties of Gases and Liquids*, 4th Edition, McGraw-Hill Book Company, 1987